

**COURSE
GUIDE**

**BIO411
PARASITOLOGY**

Course Team

Prof Chiedu Mafiana (Course Coordinator)-
NOUN
Stephen O. Omada(Course Reviwee)-



NATIONAL OPEN UNIVERSITY OF NIGERIA

© 2023 by NOUN Press
National Open University of Nigeria
Headquarters
University Village
Plot 91, Cadastral Zone
Nnamdi Azikiwe Expressway
Jabi, Abuja

Lagos Office
14/16 Ahmadu Bello Way
Victoria Island, Lagos

e-mail: centralinfo@nou.edu.ng

URL: www.nou.edu.ng

All rights reserved. No part of this book may be reproduced, in any form or by any means, without permission in writing from the publisher.

Printed 2023

ISBN: 978-058-816-X

CONTENTS

PAGE

Introduction.....iv
Course Competencies.....iv
Course Objectivesiv
Working Through this Course v
Study materials..... v
Study Units v

Introduction

BIO411: Parasitology is a 3-credit unit course in Biology. It is a 400 level, first semester undergraduate course offered to students admitted to the Faculty of Science, and Faculty of Education who are offering Biology or related programmes.

This course guide gives you a brief overview of what the course is all about, what course materials you will be using, and how you can work your way through these materials. It gives you some guidance on your Tutor- Marked Assignments.

There are In-Text Questions within each unit and Self-Assessment Exercises at the end of each unit. The exercises are overview of the unit to help you assess yourself at the end of every unit.

Course Competencies

This course aims to provide you with knowledge on concept and scope of parasitology, the ways in which parasitic diseases are spread, the different types of parasitic hosts and the types of associations that exist between parasites and their hosts.

Course Objectives

The course sets an overall objective which must be achieved through proper understanding of the course contents. Each unit has its own sets of objectives which highlight what you are expected to learn and understand. So, as you go through each unit, you should ensure that you go back to the objectives to confirm the level at which you have progressed in the course of study.

The comprehensive objectives of the course are to;

- Appreciate the various areas or aspects of parasitology
- Explain what parasites are, and their roles in the transmission of diseases
- Describe the various types of parasitic hosts
- Explain the different types of parasite – host associations
- Highlight some major groups of parasites, their taxonomy, life cycle, pathogenicity, epidemiology and management.

Working Through this Course

In this course, it is important you create enough time in reading through the course content in each unit. You would be required to do all that has been stipulated in the course: study the course units, read the recommended reference textbooks and do all the Self-Assessment Exercise(s). At some point, you may be required to submit your Tutor-Marked Assignments (TMAs) for assessment purpose. You should therefore avail yourself of the opportunity of being present during the tutorial sessions so that you would be able to compare knowledge with your colleagues.

Study materials

The course comes with a list of recommended textbooks. These textbooks are supplementary to the course materials so that you can avail yourself of reading further. Therefore, it is advised that you acquire these textbooks and read them to broaden your scope of understanding.

Study Units

This course is broken into 3 modules, with a total of fifteen units as shown below:

MODULE 1

- Unit 1 The Concept and Scope of Parasitology
- Unit 2 The Concept of Epidemiology and Parasitic Disease
Transmission
- Unit 3 Measure of Epidemiology and Disability Adjusted Life
Years DALY)
- Unit 4 Types of Animal Associations
- Unit 5 The Concept of Parasitism

MODULE 2

- Unit 1 Types of Parasitic Host
- Unit 2 Host Selection and Host Specificity
- Unit 3 Parasite and Host Associations
- Unit 4 Protozoa
- Unit 5 Trematodes

MODULE 3

- Unit 1 Cestodes

Unit 2	Nematode
Unit 3	Acanthocephalans
Unit 4	Leeches
Unit 5	Arthropods

CONTENTS **PAGE**

MODULE 1

Unit 1	The Concept and Scope of Parasitology.....
Unit 2	The Concept of Epidemiology and Parasitic Disease Transmission.....
Unit 3	Measure of Epidemiology and Disability Adjusted Life Years DALY).....
Unit 4	Types of Animal Associations.....
Unit 5	The Concept of Parasitism.....

MODULE 2.....

Unit 1	Types of Parasitic Host.....
Unit 2	Host Selection and Host Specificity.....
Unit 3	Parasite and Host Associations.....
Unit 4	Protozoa.....
Unit 5	Trematodes.....

MODULE 3.....

Unit 1	Cestodes.....
Unit 2	Nematode.....
Unit 3	Acanthocephalans.....
Unit 4	Leeches.....
Unit 5	Arthropods.....

MODULE 1

Unit 1 The concept and scope of Parasitology

- 1.0. Introduction
- 2.0. Objective
- 3.0. Content
 - 3.1. Parasitology
 - 3.2. Branches of Parasitology
 - 2.3.1 Medical Parasitology
 - 2.3.2 Structural parasitology
 - 2.3.3 Quantitative parasitology
 - 2.3.4 Parasite Ecology
 - 2.3.5 Helminthology
 - 2.3.6 Parasite Immunology
 - 2.3.7 Veterinary parasitology
 - 3.3 The scope of Parasitology
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assessment
- 7.0 References/Further reading

1.0 Introduction

Parasites are living things which harm udders by becoming metabolically dependent on them. It is an old animal relationship based on the concept of dependence for nutrition and support.

Parasitology is currently studied in a wide variety of headings and it is a dynamic aspect of zoology with links to other fields of sciences and social sciences.

2.0 Objectives

At the end of the lesson, the student should be able to clearly identify an animal that is a parasite, the branches of parasitology and its scope. Also, the student must be able to focus slides under different objectives and estimate disease prevalence

3.0 Main Content

3.1 Parasitology

No biological association is of greater medical importance than parasitism. Parasites plague humanity and are at the root cause of many diseases especially in the tropics, the need for a dedicated branch of Zoology to study these exciting but disturbing polyphyletic animal groups.

Parasitology is the study of parasites, their hosts, and the relationship between them. It is an applied field of biology dedicated to the study of the biology, ecology and relationships which parasites are involved in with other organisms known as the host.

3.2 Branches of Parasitology

Depending on the specific bias, there are different fields of parasitology and some of these include medical parasitology, veterinary parasitology, structural parasitology, quantitative parasitology, parasite ecology, conservation of parasites, malariology, helminthology parasite immunology etc.

3.2.1 Medical Parasitology

This is the science that deals with organisms living in the human body (the host) and the medical significance of host-parasite relationship. It's also concerned with the various methods of their diagnosis, treatment and finally their prevention and control.

3.2.2 Veterinary parasitology

The study of parasites that cause economic losses in agriculture or aquaculture operations, or which infect companion animals. This is becoming particularly important as emerging diseases threatens global food security. In Nigeria, a number of studies have been conducted in fish parasitology and other vertebrates such as Lizards and Cats with Zoonotic potential.

3.2.3 Structural parasitology

This is the study of structures of proteins from parasites. Determination of parasitic protein structures may help to better understand how these proteins function differently from homologous proteins in humans. In addition, protein structures may inform the process of drug discovery.

3.2.4 Quantitative parasitology

Parasites exhibit an aggregated distribution among host individuals, thus the majority of parasites live in the minority of hosts. This feature forces parasitologists to use advanced biostatistical methodologies.

3.2.5 Parasite Ecology

Parasites can provide information about host population ecology. In fisheries biology, for example, parasite communities can be used to distinguish distinct populations of the same fish species co-inhabiting a region. Additionally, parasites possess a variety of specialized traits and life-history strategies that enable them to colonize hosts. Understanding these aspects of parasite ecology, of interest in their own right, can illuminate parasite-avoidance strategies employed by host

3.2.6 Malariology

3.2.7

This is an aspect of parasitology which focuses mainly on the study of Protozoan parasite, *Plasmodium*, its species, their biology, pathogenicity, epidemiology and management of the parasitic infection.

3.2.8 Helminthology

3.2.9

“Helminth” means worms. Hence as the name implies it is the study of vermiform parasites ranging from trematodes to Cestodans, Nematodans and leeches.

3.2.10 Parasite Immunology

This is an aspect which deals with parasite survival in host as well as host susceptibility. This aspect is particularly important when formulating concentrations of chemotherapeutic agents and vaccines.

In-Text Question

- **What are parasites?**

Answer

- **Parasites are living things which harm udders by becoming metabolically dependent on them**
-

Self-Assessment Exercise 1

- **Define parasitology?**
- **List 5 branches of parasitology**

3.3 The scope of Parasitology

As a biological discipline, the scope of parasitology is not determined by the organism or environment in question, but by their way of life. This means it forms a synthesis of other disciplines, and draws on techniques from fields such as cell biology, bioinformatics, biochemistry, molecular biology, immunology, genetics, evolution and ecology.

Thus, parasitology covers a wide range of topics which center around parasites, their taxonomic positions, their structural and physiological biology, life cycle patterns, ecology, pathogenicity, diagnosis of

infection as well as management with chemotherapy and other contemporary techniques.

Self-Assessment Exercise 2

- **What is the scope of parasitology?**

4.0 Conclusion

Parasitology has become an established branch of Applied Zoology which has been able to integrate with other branches of sciences as well as social sciences in managing parasitic diseases world over.

5.0 Summary

- Parasites are a polyphyletic group of animals ranging from protozoans to chordates
- They are responsible for a number of diseases and mortality in their hosts which includes humans.
- In order to properly reflect the complex interactions of parasites and their host, they are studied under diverse branches.

7.0 Tutor Marked Exam

- a. Define parasitology and write short notes on any five branches of parasitology

7.0 References/Further reading

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp

Ukoli, F.M.A. (1990). Introduction to parasitology in Tropical Africa. Texflow Limited, Ibadan. 462pp.

Answers to SAEs

SAE 1

- Parasitology is the study of parasites, their hosts, and the relationship between them.
- Branches of parasitology include; Medical parasitology, Helminthology, Parasite ecology, Medical parasitology and Parasite immunology.

SAE 2

Scope of parasitology; parasitology covers a wide range of topics which center around parasites, their taxonomic positions, their structural and physiological biology, life cycle patterns, ecology, pathogenicity, diagnosis of infection as well as management with chemotherapy and other contemporary techniques.

Unit 2 The concept of epidemiology and Parasitic Disease Transmission

Content

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1. Parasitic Disease transmission
 - 3.2. The concept of epidemiology
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
 - a. Draw the disease triad and use it in explaining parasitic disease transmission
 - b. Design an epidemiological methodology for assessing malaria diseaseprevalence.
- 7.0 References/Further reading

1.0 Introduction

Parasites are dynamic in their distribution-some are endemic while many are ubiquitous. The environment plays a key role in their survival and transmission. Often times, they rely on vectors (mostly arthropods) for their transmission.

2.0 Objective

At the end of the class, student must have fully understood, the distribution of parasites and their general effects on biological systems and disease transmission patterns. The student should also be able calculate the prevalence of diseases.

3.0 Main content

- 3.1. Parasitic Disease transmission
- 3.2. The concept of epidemiology


3.1 Parasitic Disease transmission

By virtue of their nature, parasites reside in their hosts, depend on their hosts for physiological and nutritional ends and hence often result in adverse effects on their host.

The parasite-host interaction does not occur in isolation; rather it is affected by conditions known as risk factors. A risk factor which must occur for a disease situation to arise is known as the necessary risk while the organism/parasite which causes the disease is called an agent.

The causal web is a simple triangle which shows the inevitable link

between the agent (parasite), the host and the environment. Often times especially with parasitic infections, there may be a vector which links all components as shown below.



Thus, a number of factors predisposes a host to attack by a vector, affects vector viability and life cycles and even how the host respond to parasitic infections/diseases. All these add up into the concept of Epidemiology.

In-Text Question

- **Parasites rely mostly on _____ for their transmission**

Answer

- **Vectors**

3.2 The concept of epidemiology

Epidemiology is the study of the disease pattern and distribution, within a defined community or population.

It is derived from the combination of Greek words Epi which means upon, demos which means people and logy which means study.

It is an interdisciplinary subject, involving medicine, science, geography and statistics. Epidemiology is a quantitative science that evaluates the occurrence, distribution, determinants of control of disease pattern in a defined population. Thus, in epidemiology, demographic and social characteristics of the host such as population density, poverty etc. are important attributes in epidemiological studies. Physical, biological, sociological and psychological variables are significant determinants of disease distribution and patterns. They are considered as extrinsic factors. Demographic and social attributes of the definitive host are significant factors in disease manifestation. The demographic factors include:

- Age

- Sex,
- Race,
- Population density etc.

Thus, epidemiology is a scientific enquiry on disease occurrence. i.e., the study of disease distribution and its determinants. The unit of the study of epidemiology is a defined or identifiable population. A population in this context is not always defined by geographical boundaries rather by a number of other factors such as age, sex, occupation, ethnicity, religion, institution, etc.

Epidemiology is therefore the study of disease occurrence in populations- humans, plants, animals.

It is indeed an analytical arm of community and preventive medicine, utilizing statistics as a working tool. It is integral to the management of public health and it aims to improve the health of the population.

The aims of epidemiology include to:

- Determine the primary agent or ascertain causative factors
- Understand the causation of disease, disorders or conditions
- Determine the characteristics of the agent or causative factors
- Define the mode of transmission
- Define and determine contributing/confounding factors
- Identify and explain geographic disease patterns
- Determine, describe and report the natural cause of disease, disability, injury and/or death
- Determine suitable control measures
- Determine preventive measures
- Aid in planning and development of health services and
- Provide administrative and planning data in public health management.

Self-Assessment Exercise

- **Define epidemiology**
- **Mention 4 factors that are significant in disease manifestation**
- **List 5 aims of epidemiology**

4.0 Conclusion

The parasitic disease triad incorporates the agent (parasite), host and the environment as the key determinants of parasitic disease transmission.

5.0 Tutor Marked Assignment (TMA)

- a. Draw the disease triad and use it in explaining parasitic disease transmission
- b. Design an epidemiological methodology for assessing malaria disease prevalence.

6.0 References/Further reading

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp.

Answer to SAE

1. Epidemiology is the study of the disease pattern and distribution, within a defined community or population.
2. Factors that are significant in disease manifestation are;
 - a. Age
 - b. Race
 - c. Population density
 - d. Sex
3. Aims of epidemiology include to;
 - a. Determine suitable control measures
 - b. Determine preventive measures
 - c. Aid in planning and development of health services and
 - d. Provide administrative and planning data in public health management
 - e. Define the mode of transmission

Module 1

Unit 3 Measure of Epidemiology and Disability Adjusted Life Years (DALY)

Content

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Measure of epidemiology
 - 3.2 Disability Adjusted Life Years (DALYs)
 - 3.2.1 DALYs estimate of Common Parasitic Diseases
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/Further reading

1.0 Introduction

Epidemiological estimates have become important at local, national and international scales for the provision of health statistics for the purpose of budgeting and forecasting. Hence a set of statistical indices have been evolved to meet this end. This has been coupled with new innovative techniques which involves the use of Rapid assessments (e.g RAPLOA, REMO), Remote sensing (RS) and Geographic Information Systems (GIS).

2.0 Objectives

At the end of the class, student must be familiar with the terms Prevalence, incidence, intensity, disease burden and DALYs, DALY estimates for different parasitic diseases and how to estimate them.

3.0 Main Content

3.1 Measure of Epidemiology

There are a number of tools for measurement or description in epidemiology. These include:

- Prevalence: frequency of a disease at a particular time. Based on prevalence, a disease may be defined as endemic (maintain a relatively constant but moderate level of occurrence), Hyper- endemic (persistently high and continuous level of occurrence), epidemic (an outbreak or sudden occurrence of a disease with considerable intensity) or Holoendemic (high prevalence amongst children in a population).
- Epidemiological rate- morbidity rate, mortality rate, incidence rate, admission rate etc.
- Disease Burden (overall weight of effect of a disease) using DALY.

- Estimation of disease intensity using direct parasite count, indirect counting of eggs and cysts in faeces, indirect measurement of levels of immunological responses to parasite antigens.
- Parasite dispersion estimation which may be under dispersion, random or over dispersion.
- Geographical distribution estimation using mapping techniques such as annual cartographic maps, geographic information system and remote sensing.

Epidemiological studies may be descriptive, analytical (cohort, or case control), experimental or rapid epidemiological assessment.

In-Text Question

- Differentiate between endemic and hyper-endemic prevalence of a disease

Answer

- An endemic disease maintains a relatively constant but moderate level of occurrence while a hyper-endemic disease is persistently high with a continuous level of occurrence.

3.2 Disability Adjusted Life Years (DALYs)

In epidemiology, the disease burden on the economy is approached from the view that morbidity from parasitic infection impairs productivity as a result of loss of valuable man- hour. This is calculated using an index known as the Disability adjusted life years (DALYs).

DALYs is a health indicator which translates disabilities into years of healthy life lost, by giving each disease state a disability weight ranging from zero (healthy) to 1 (death). It incorporates the potential years of lost life as a result of death at a given age. It measures the burden of a disease of a defined population and the effectiveness of interventions. They are valid indicators of population health.

In summary, DALYs can simply be defined as the number of health years of life lost due to premature death or disability.

3.2.1 DALYs estimate of Common Parasitic Diseases

The table below shows the global estimates of parasitic disease effects using the DALYs tool. Table 1: Global estimates of some parasitic diseases (WHO, 2002)

Disease	TDR Diseases	DALYs (hundreds of thousands)	Deaths of (Thousands)	Pop. At risk of mill	No. of endemic mic	No. of

	Category	Total	Male	female	Total	Male	female	ons	count	infected prevalence
African trypanosomiasis	1	1,598	1029	568	50	32	18	50m	36	>300,000
Leishhmaniasis	1	2357	1410	946	59	35	24	350	82	12m
								m		
Malaria	2	42280	20024	22256	1124	532	592	200m	90	300-500m

The burden of various diseases over time, depends on many factors. Presently, some disease burdens have continued to increase. For instance, increasing malaria burdens occur in some countries of Africa due to factors such as drug resistance, frequent exposure of the non-immune populations, the emergence of HIV/AIDS, climatic and environmental changes and the breakdown of control programmes. Such increased parasite burden results in increased economic burden of the population.

The economic burden of a disease is the totality of direct and indirect costs associated with the infection. These include morbidity, loss of productivity, absenteeism and cost of health care in the infected people, as well as the losses in agriculture and non- agricultural enterprises.

Self-Assessment Exercise

- Define prevalence
- Briefly explain Disability Adjusted Life Years

4.0 Conclusion

Given the economic burden associated with diseases and death resulting from parasitic diseases, it becomes important to have a well developed scheme to attributing relative importance to diseases and resource allocation therefore.

Physical, biological, sociological and psychological variables are significant determinants of disease distribution and patterns.

Epidemiology incorporates to a large extent mathematical and geographical tools in solving parasitic disease related public health issues.

5.0 Summary

Epidemiological tools help in explaining parasitic disease problems by incorporating geographical, temporal and mathematical aspects.

6.0 Tutor Marked Assignment (TMA)

- a. Justify the use of DALYs in parasitic disease management programs
- b. State the relative importance of different parasitic diseases on the basis of their DALYs estimate.
- c.

7.0 References/Further reading

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp.

Answer to SAE

- Prevalence is defined as the frequency of a disease at a particular time. Based on prevalence, a disease may be defined as endemic, Hyper-endemic, epidemic or Holoendemic.
- DALYs is a health indicator which translates disabilities into years of healthy life lost, by giving each disease state a disability weight ranging from zero (healthy) to 1 (death). It incorporates the potential years of lost life as a result of death at a given age. It measures the burden of a disease of a defined population and the effectiveness of interventions. They are valid indicators of population health.

Unit 4 Types of animal associations

Content

- 1.0 Introduction
- 2.0. Objective
- 3.0. Main content
 - 3.1 Homospecific/intraspecific/ homogenetic and
 - 3.2 Heterospecific/interspecific/heterogametic
 - 3.3 Loose associations
 - 3.4 Temporary associations
 - 3.5 Intimate associations
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further reading

Organisms live within various habitats ranging from air, water and soil; interacting with each other in the environment. These associations are characterized by diverse benefits: provision of food, transport, protection, etc.

Within the environment, living organisms exhibit diverse forms of associations which may be broadly divided to two categories namely:

- Homospecific/intraspecific/ homogenetic and
- Heterospecific/interspecific/heterogametic

2.0. Objective

Proper understanding of animal association, with examples of host-parasite interactions and the challenges therein. Ability to carry out basic sample preparation for microscopy and common parasite identification.

3.0. Main Content

3.1 Homospecific/intraspecific/ homogenetic

This deals with relationships between organisms/members of the same species. E.g associations such as a flock of sheep, a school of fish, herds of cattle, colonies of coelenterates, filaments of algae, colonies of social insects (such as bees, termites and ants) etc. These associations are based on cooperative behaviors within the groups. However, this association falls outside the scope of parasitism.

3.2 Heterospecific/interspecific/heterogametic

This occurs between organism of different species, resulting from diverse biological links in morphology, ecology, physiology and biochemistry.

A spectrum of heterospecific associations may overlap, consisting of predation, phoresis, commensalism and parasitism.

Predation is non-specific and usually results in the death or injury of one of the interacting organisms. The degree of predation may be absolute, partial or temporary or insignificant. It forms the basis of interspecific associations among living things. Heterospecific relationships are collectively called symbiosis. Animals may exhibit different types of symbiotic relationships which may be obligate or facultative. The three basic types of heterospecific relationships are:

- **Loose associations**
- **Temporary associations**
- **Intimate associations**

In-Text Question

- A school of fish is an example of what category of association?
-

Answer

- **Homospecific/intraspecific**

3.3 Loose Associations

This form of association is casual because they retain their physiological independence and are just ecologically linked. Loose associations provide biological advantages to those organisms in the association over those that are not in association. Significantly, it is usually one-sided benefit. They include:

Phoresis Commensalism

Phoresis (transport commensalism) is a form of loose association in which an organism provides shelter, support or transport for the other organism, usually a sedentary animal of a different species. The passenger can survive without its usually motile substratum or attachment surface. There is no conflict of interest in this association. Examples are barnacles on limpet mollusks, dung beetle, *Aphodius fimentarius* carrying dauer-larvae of *Pledora coactatata* (a free-living nematode which feeds on bacteria) etc.

Commensalism (Co= together, Mensa= table) is an association between two organisms in which one benefits, but the host is not hurt nor injured. It usually occurs between species that are either vulnerable to predation, danger or with an inefficient means of locomotion, feeding or defence. It is the most indefinite and least obligatory inter-specific relationship. The host neither benefits nor loses in the relationship.

There are three types of commensalism

- a. **Cleaning:** in which one organism cleans the other by feeding on

the parasite of the host organism. E.g *Bulbus ibis* (cattle egret) feeds on ectoparasites on the skin of cattle, *Naucrates doctor* (pilot fish) feeds on the ectoparasites of shark.

b. Protection/camouflage. E.g the anemone hermit crab, *Dardanus pedunculatus* attaches several anemones to its shell, for camouflage and as a deterrent to possible predators.

c. Synoecious (syn= together, oikos= house): which involves animal living together in the shell/burrow of another animal. E.g *Mytilus edulis* (mussel) harbors the pea crab (*Pinnotheras pisum*) within its mantle cavity.

In-Text Question

- **An association between two organisms in which one benefits, but the host is not hurt nor injured is referred to as?**

Answer

- **Commensalism**

3.4 Temporary associations

This form of association is typified by the predator- prey relationships in which one organism uses the other as a source of food. The predator usually kills the prey or eats part of it (as in plants). E.g leeches feeding on blood of aquatic animals and some blood sucking insects such as lice and ticks.

3.5 Intimate associations

There are two basic forms of intimate associations. They are:

- **Mutualism**

This is a non-competitive, physiologically interdependent association between organisms of two different species. It is considered as the most advanced form of symbiosis because it involves reciprocal benefits. They always live together and mutualism is recognized as a special case of parasitism, in which metabolic by-products of the associates are of value to the partners. E.g *Trichonympha campanula* (intestinal flagellates) feeds on the wood chips in the gut of termites and wood roaches. The by-products of cellulose digestion and cellulose of the flagellated protozoan are useful to the insect. The sexual reproduction of the protozoa is stimulated and regulated by the insect moulting hormone, ecdysone. The anaerobic environment of the insect's gut is integral to the survival of the flagellate. The two organisms depend on each other for survival. Physiological dependence is an essential component of mutualistic relationships.

4.0 Conclusion

Parasite association with the host confers them so much advantages for survival, hence the ultimate aim of parasitism unlike predation is not the

mortality of the host, rather a parasite requires its host to survive to ensure continued supply of food materials.

5.0 Summary

- Parasite-host relationship is a form of symbiosis
- The parasite benefits at the expense of the host
- The relationship may be loose, temporary or intimate

6.0 Tutor-marked Assignment

State the different kinds of parasite-host associations and describe the parasite adaptations for each

7.0 References/Further reading

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp.

Module 1

Unit 5: The concept of Parasitism

- 1.0** Introduction
- 2.0** Objective
- 3.0** Main content
- 3.1** Parasitism
- 3.2** Types of parasites
- 4.0** Conclusion
- 5.0** Summary
- 6.0** Tutor Marked Assignment (TMA)
- 8.0** References/Further reading

Content

1.0 Introduction

Parasitism (para= beside; sitos=food).

Parasitism is at the root of several diseases and deformities in many vertebrates. The degree of effect depends on the type of parasite, its mode of action and of course the host physiological state.

2.0 Objective

At the end of the class, student must be familiar with the types of parasites and hosts and the basis for parasite and host categorization.

3.0 Main content

3.1 Parasitism

This is a physiological association in which one organism (the parasite) benefits while the host loses. It is one-sided and it is a social and ecological relationship between two different organisms, one being injured in the association.

Parasitology therefore is defined as an aspect of symbiology which deals with the study of parasites. It is a medical and ecological science that encompasses the study of parasite and parasitic diseases.

A parasite is defined as an organism which has a detrimental effect on the intrinsic growth rate of its host population

Parasitism is also defined as an association, generally continuous, between two different organisms, one of which lives at the expense of the other.

In-Text Question

- **Differentiate between parasite and parasitism**

Answer

- **A parasite is an organism which has a detrimental effect on the intrinsic growth rate of its host population, while parasitism is**

an association, generally continuous, between two different organisms, one of which lives at the expense of the other.

3.2 Types of parasites

Parasites may be plants, animals, bacteria or viruses and they occur in every phylum in the Kingdom animalia, ranging from protozoans to chordates. They are categorized based on different criteria.

3.2.1 Site of residence or habitat:

Ectoparasites Endoparasites

3.2.2 Duration or degree of dependence

Temporary parasites Permanent parasites

3.2.3 Nature of host

Hyper-parasites (parasites of parasites)

Cleptoparasites (organisms that steal food reserves of its host to feed its young)

3. 2.4 Other types of parasites includes Wandering /Errant/Aberrant parasites Occasional/Accidental parasites Obligatory/Obligat parasites Facultative parasites

Microparasites Macroparasites Pseudoparasites

Coprozoic or spurious parasite

4.0 Conclusion

Parasites vary in taxa, mode of action, host selection and behavior and these factors together determines the pathogenicity and hence the importance attributed to a given parasite.

5.0 Tutor Marked Assignment (TMA)

- a. what is a parasite?
- b. state the criteria for categorizing parasites.

6.0 References/Further reading

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp.

MODULE 2:

Unit 1: Types of Parasite Host

Content

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
- 3.1 Types of Hosts
- 3.2 Host-parasite interaction
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/further readings

1.0 Introduction

The Host is a key component of parasitic relationship and some parasite may require more than one host to complete their life cycle. Others simply have alternative host, infecting a broad range of animals.

2.0 Objective

At the end of the class, student must be familiar with the types of hosts and the basis for parasite and host categorization.

3.0 Main content

3.1 Types of hosts

A host is the non-parasitic member in a parasitic association. It harbors the parasites and thus provides itself as the environment of the parasite, albeit involuntarily. Biotic and abiotic factors of the host influence the survival, distribution, prevalence, intensity, etc. of the parasite. Abiotic factors in the host include:

- pH
- O₂
- CO₂

There are different types of hosts, depending on a number of criteria. They include:

Definitive or Primary host
Intermediate or secondary host
Reservoir host
Paratenic/transport/carrier host
Ecological host
Vector
Dead end host

In-Text Question

- _____ is the non-parasitic member in a parasitic association

Answer

- **Host**

Hosts, parasite stage and host necessity

Type of host	Stage of parasite in the host	Necessity of the host
Definitive	Adult	Essential
Intermediate	Juvenile	Essential
Paratenic	Juvenile	Ecologically necessary
Vector	Juvenile	Essential
Reservoir	Adult	Not Essential

3.2 Host-parasite Interactions/Relationships

Generally, parasites may locate their host directly or indirectly and then establish a relationship that is determined by ecological, physiological, geographical and evolutionary factors. The selective and dynamic phenomenon of parasite adaptation to certain hosts is known as host specificity.

Specificity results from a number of convergent phenomena, starting with habitats, microhabitat and feeding styles of the free-living progenitors. This influences the types of interaction parasites have with the host and through this specificity results.

In-Text Question

- **Define host specificity.**

Answer

- **Host specificity is the selective and dynamic phenomenon of parasite adaptation to certain hosts**

4.0 Conclusion

Parasite-host interaction is dynamic and is intimately tied to their life cycle selection.

5.0 Summary

The different categories of parasite hosts are definitive, intermediate, paratenic, vector and reservoir.

Depending on the host different life cycle stages of the parasite may be found and some hosts may not be essential in the life cycle

6.0 Tutor Marked Assignment (TMA)

- a. What is a host?
- b. What are the host abiotic factors which form the parasite ecosystem?

7.0 References/further readings

Yoloye, L.V. (1984) *Invertebrate Zoology*. Ilorin University Press, Ilorin. 318pp.

Otubanjo, O.A. (2008) *Elements of Parasitology* Second Edition. Panaf Publishing Inc. Abuja 196pp

Unit 2 Host Selection and Host specificity

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Host Selection
 - 3.2 Host specificity
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/Further reading

1.0 Introduction

Parasite-Host interaction is often specific, with a parasite living on a single host or on closely related host taxa. Their choice of host and host specificity depends on a number of factors ranging from structural, physiological to environmental.

2.0 Objective

At the end of the class, student must be familiar with the different factors which influence parasite host selection, specificity, adaptations and co-evolution with its host as well as immunological factors leading to host resistance and host protection.

3.0 Main content

3.1 Host Selection

Parasites may possess single or a wide range of hosts in their life cycle. They may be:

Monoxenous i.e., if they possess a single host, **Oligoxenous** i.e., if they utilize a small range of host **Polyxenous**, i.e., if they use many suitable hosts.

Parasites with indirect life cycles are less host specific than those with direct life cycles. Thus, there is less host specificity with increase in the number of intermediate hosts. Co-evolutionary factors are important in parasites that feed on host tissues or occur in extra intestinal sites. This is because the parasite must present the right cues/signals and respond appropriately to host defense system. The inability of a parasite to respond to such cues may act as a barrier on its host range.

In-Text Question

- **Parasites that utilize a small range of hosts are said to be?**

Answer

- **Oligoxenous**

3.2 Host specificity

The host range or host specificity of a parasite is the collection of hosts that an organism can utilize as a partner. In the case of human parasites, the host range influences the epidemiology of the parasitism or disease. Specificity in parasites is seldom absolute. It differs amongst various types of parasites and at different stages of the parasite life in the definitive and intermediate hosts. Majority of parasites however are restricted to just one or few closely related species.

Supra –specific condition occurs when groups of parasites are associated with natural groups of hosts.

Infra-specific conditions on the other hand occur when a single species of parasite is associated with a single host. Over 70% of described monogenetic trematodes are infra- specific.

Phylogenetic, physiological, ecological and human factors affect parasite specificity. Specificity is the effect of biological compatibilities between hosts and parasites. The range of host can be limited by lack of host stimuli, host immune response, innate resistance and simply the availability of essential nutrients to the parasite.

Localization of, establishment, growth and reproduction are factors which contribute to host specificity. Behavioral adaptations have also contributed tremendously to increase the chances of host location. Chemical cues emitted by hosts play vital roles in the infective stages locating or finding their hosts. Host selection is determined by morphological and physico-chemical factors.

In-Text Question

- _____ condition occurs when groups of parasites are associated with natural groups of hosts.

Answer

- **Supra-specific**

4.0 Conclusion

The success of parasitism depends on the specificity of host parasite relationships. Ability to secure the appropriate host is key to parasite survival and hence it attracts a lot of energy investment, often requiring co-evolution with the host.

5.0 Tutor Marked Assignment (TMA)

- What is host specificity?
- list the criteria for parasite host selection

6.0 References/Further reading

Croll, N.A. (1973). Parasitism and other associations. Pitman medical, London, 98pp.

Otubanjo, O.A. (2008) *Elements of Parasitology* Second Edition. Panaf Publishing Inc. Abuja 196pp

Module 2

Unit 3 Parasite and Host Associations

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Parasite adaptations and evolution
 - 3.2 Host resistance and Host protection
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 6.0 References/Further reading

1.0 Introduction

Host specificity is in a state of constant flux and is a product of evolution. Host specificity, maybe lost in the course of evolution, thus reflecting the dynamism of parasite host relationships.

2.0 Objective

At the end of the class, student must be familiar with the different factors which influence parasite host selection, specificity, adaptations and co-evolution with its host as well as immunological factors leading to host resistance and host protection.

3.0 Main content

3.1 Parasite adaptations and evolution

Host–parasite co-evolution is a special case of co-evolution, which is defined as the reciprocal adaptive genetic change of two antagonists (e.g., different species or genes) through reciprocal selective pressures. In the particular case of host–parasite co-evolution the antagonists are different species of host and parasite. Hosts and parasites exert reciprocal selective pressures on each other, which may lead to rapid reciprocal adaptation. For organisms with short generation times host–parasite co-evolution can be observed in comparatively small time periods, making it possible to study evolutionary change in real-time under both field and laboratory conditions.

Specificity patterns are shaped during the course of parasite evolution. Phylogenetic specificity is determined by ecological or physiological factors or a combination of both.

The evolutionary basis of host specificity is supported by the fact that primitive hosts harbor primitive parasites. Hosts of monophyletic (single) are parasitized by an exclusive group of parasites. Parasites that

evolved with their hosts are known as heirlooms while those that are picked up through ecological links or contact are known as souvenirs. E.g., adults of family Taenidae parasitize only mammalian hosts. Elasmobranchs are exclusive hosts of the tapeworms of the order Tetrathyridida and tetrahymincha.

In-Text Question

- **What is the difference between heirlooms and souvenirs?**

Answer

- **Heirlooms are parasites that evolved with their hosts, while souvenirs are parasites picked up through ecological links or contact.**

3.2 Host resistance and Host protection

Free-living organisms struggle predominantly with abiotic factors within the environment for their survival. Biotic and abiotic factors of parasite habitats affect the establishment, maintenance and survival in the host.

They experience varied destructive factors in their host and their ability to establish, survive and maintain their presence in their host is always under threat. They are affected by host immunity which constantly produce antibody against parasite.

Host resistance refers to the unsuitability of a host. This is however never absolute. It may result from the presence of some physical or chemical barriers that prevents parasite from establishing in a host. A number of host characteristics such as physiology, behavior and structure influence host-parasite interactions, thereby influencing host resistance. Host natural resistance can also arise through geographical distribution, behavioral characteristics of the host or nutritional habits as the host may not come in contact with the infective stages.

Natural resistance encompasses a physiological incompatibility between parasite and host, preventing parasite invasion, establishment and survival, without the intervention of immunological- based protective responses. It is influenced by factors such as host's genetic composition, age, nutritional status and gender.

Host protection

Organisms are able to distinguish between self and non-self using immune response. While many parasites have evolved successful means of evading host immune response, the ability of the host to control the parasite is not absolute. Various host exhibit different degrees of susceptibility or resistance for parasitic agents or specific infections.

Hosts are protected from infection by:

- 4 Non-specific resistance and
- 5 Specific immunity or protection

Non-specific resistance is also known as innate or natural immunity. It

includes mechanisms which the host is genetically endowed with at birth and thereby protected from certain infections.

4.0 Conclusion

Parasitism requires high energy investment on the part of the parasite to evade host immunological responses.

5.0 Summary

- Parasites have to adapt to host conditions to survive.
- Hosts have innate mechanism to fight against parasites.
- The success of parasitism depends on the parasites ability to evade host immuneresponse.

6.0 Tutor Marked Assignment (TMA)

- a. List the factors that affect host-parasite co-evolution
- b. write short notes on host resistance and host protection

7.0 References/Further reading

Otubanjo, O.A. (2008). Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp.

Unit 4 Protozoa

Content

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Parasitic Protozoology
 - 3.4 Life Cycle
 - 3.5 Pathogenicity
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor- Marked Assignment
- 7.0 References/Further reading

1.0 Introduction

Protozoans are a polyphyletic group of unicellular animals whose mode of classification is primarily based on their locomotory organelle. All protozoan phyla contain parasitic members. The mode of transmission, location within host and pathogenicity varies widely. Protozoans are the most common parasites of vertebrates.

2.0 Objective

Students would at the end of this lesson, be conversant with the different types of protozoan parasites, their taxonomy, characteristics, life cycle, pathogenicity, epidemiology and management. Basic laboratory parasite assessment skill using light microscopy.

3.0 Main content

3.1 Taxonomy

The basic taxonomic character for this group is the absence/presence and type of motile organelles

Sarcodina e.g. *Entamoeba*

Mastigophora e.g. *Trypanosoma*, *Trichomonas*, *Leishmania*, *Opalina*, *Giardia*

Ciliophorans e.g. *Balantidium*

Apicomplexa e.g. *Toxoplasma*, *Plasmodium*, and *Eimeria*.

In-Text Question

- _____ are the most common parasites of vertebrates.

Answer

- **Protozoans**

3.2 Biology (Characteristic Features)


Protozoans are generally unicellular, microscopic eukaryotic organisms. They are the first animals with membrane-bound nucleus, as well as heterotrophism. A considerable number of these organisms have motile organelles; while majority have at least one stage of their life cycle being motile.

3.3 Parasitic Protozoology (Protozoans of parasitic importance)

These include *Entamoeba spp* (*E. histolytica*, *E. coli*, *E. gingivalis*), *Opalina ranarum*, *Trypanosoma* (*T. brucei brucei*, *T. brucei rhodesiense*, *T. brucei gambiense*, and *T. cruzi*), *Trichomonas* (*T. vaginalis*), *Leishmania* (*L. donovani*, *L. aethiopica*), *Giardia lamblia*, *Balantidium coli*, *Toxoplasma gondii*, *Plasmodium* (*P. falciparum*, *malariae*, *ovale*, and *vivax*), and *Eimeria*.

3.4 Life Cycle

This is generally asexual, involving binary fission, multiple fissions (schizogony, Merogony, Sporogony, Polyembryony), budding, and sexual (syngamy, autogamy, and conjugation). Many of these organisms have direct life cycle (wherein parasites undergo complete development within the host), while some others have part of their life cycle completed outside the host; this could be a substratum, intermediate host and/or vectors.



Life cycle of Protozoan parasite (Plasmodium sp.) which causes malaria (Centre For Disease Control, USA).

In-Text Question

- Mention 3 protozoans of parasitic importance

Answer

- *Entamoeba spp*, *Trichomonas (T. vaginalis)*, and *Giardia lamblia*.

3.5 Pathogenicity

Many of these parasites are found in nutrient-rich sites such as the GIT (especially the small intestine) liver, blood plasma, blood cells, and other tissues Epidemiology, Control and Management.

4.0 Conclusion

Protozoans are a polyphletic group of unicellular animals. There are parasitic protozoan species in all the taxonomic classes. They are ubiquitous and are among the most common and most deadly parasites.

5.0 Tutor- Marked Assignment

a. Draw, label and write short notes on 5 protozoan parasites b. Describe the Life cycle of the malaria parasite (*Plasmodium* spp)

6.0 References/Further reading

Yoloye, L.V. (1984) Invertebrate Zoology. Ilorin University Press, Ilorin. 318pp.

Otubanjo, O.A. (2008) Elements of Parasitology Second Edition. Panaf Publishing Inc. Abuja 196pp

Ukoli, F.M.A. (1990). Introduction to parasitology in Tropical Africa. Texflow Limited, Ibadan. 462pp.

Unit 5 Trematodes

Content

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Life Cycle
 - 3.4 Pathogenicity
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked assignment
- 7.0 References/Further reading

1.0 Introduction

Trematodes constitute a group of parasitic flat worms commonly referred to as flukes. They may be Monogenetic or digenetic-The latter are characterized by two hosts, a definitive host and an intermediate host usually a mollusc.

2.0 Objective

Students would at the end of this lesson, be conversant with the different types of trematodes of parasitic importance, their taxonomy, characteristics, life cycle, pathenogenicity, epidemiology and management. Preparation

3.0 Main content

3.1 General Taxonomy of Helminthes

Flatworms is broadly divided into two classes; Monogeneans and digeneans. The monogenetic trematodes are so called due to the completion and dependence on a single host, and are common with aquatic vertebrates (fishes). Examples are *Dactylogyrus* and *Gyrodactylus*. Digenetic trematodes compulsorily complete their life cycle between two different host species. The most common ones include *Fasciola* (*F. hepatica* and *F. gigantica*), *Schistosoma* (*S. haematobium*, *S. mansoni*, *S. japonicum*, *S. intercalatum*, *S. mekongi*), *Paragonimus westermani*, and *Clonorchis sinensis*.




Figure 1; Life Cycle of the Trematode (*Schistosoma* sp)

In-Text Question

- Trematodes that are characterized by two hosts; a definitive host and an intermediate host, are referred to as

Answer

- Digenetic

3.2 Biology (Characteristic Features)

These are bilaterally symmetrical triploblastic organisms, pronounced cephalization, and the onset of specific physiologic organs. However, parasitic flatworms are devoid of respiratory, circulatory, and digestive systems absent. Trematodes are generally flat, elongate and non-segmented worms.

3.3 Life Cycle:

Parasitic flatworms are basically hermaphroditic (except *Schistosoma spp*)

3.4 Pathogenicity

These parasites invade organs such as musculature, lungs, renal and intestinal blood vessels, as well as liver of mammals.

3.5 Epidemiology and management

Trematode infection is common in the tropic particularly in places where there is a preponderance of their snail intermediate host.

4.0 Conclusion

Trematodes constitute an important class of parasites with both human and a veterinary burden.

5.0 Summary

- Trematodes are often referred to as flukes
- Common flukes include blood fluke (*Schistosoma* sp.), liver fluke (*Fasciola* sp.), lung fluke (*Paragonimus* sp.)
- They are digenetic, requiring an intermediate host, mostly mollusk for their transmission.

6.0 Tutor Marked Assignment

- a. List the common Digenetic trematodes, their common and scientific name as well as their hosts.
- b. Write short note on schistosomiasis

7.0 References/Further reading

- Yoloye, L.V. (1984) *Invertebrate Zoology*. Ilorin University Press, Ilorin. 318pp.
- Otubanjo, O.A. (2008) *Elements of Parasitology* Second Edition. Panaf Publishing Inc. Abuja 196pp
- Hickman *et al.* (2008). *Integrated principles of Zoology*. 14th Edition, Mc-Graw Hill. 910pp

MODULE 3

Unit 1 Cestodes

- 1.0 Introduction
- 2.0. Objective
- 3.0. Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.2 Life Cycle
 - 3.4 Pathogenicity
 - 3.5 Epidemiology and Management
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor- Marked Assignment
- 7.0 References/Further reading

1.0 Introduction:

Cestodes are flatworms, commonly referred to as tapeworms because of their long tape-like appearance, with some growing up to several meters long. They are endoparasites of vertebrates such as fishes, swine, cattle, cats, man and a host of others.

2.0. Objective

At the end of the class, student must be familiar with the different species of cestodes, their host, mode of transmission and management practices.

3.0. Main content

3.1 Taxonomy:

Members are of the Class Cestoda, Sub-Orders Eucestoda and Cestodaria. Examples include *Taenia* (*T. solium*, *T. saginata*), *Diphyllobothrium latum*, *Dipylidium caninum*, *Hymenolepis nana*, and *Echinococcus granulosus*

3.2 Biology (Characteristic Features)


These are flat, elongated and segmented worms. These segments progressively increase in size and complexity away from the cephalic regions (Strobilization). Immature proglottids are closer to the head, mature ones intermediate, while gravid ones are farthest from the head. Head region is provided with adhesive suckers and hooks for attachment to host for phoretic and nutritional purposes.

In-Text Question

- Give 3 examples of cestodes

Answer

- *Taenia* (*T. solium*), *Echinococcus granulosus*, and *Diphyllobothrium latum*.



A typical Tapeworm (*Taenia* sp.)
 (http://www.medicalook.com/diseases_images/tapeworm2.gif)

3.3 Life Cycle

Members are all hermaphrodites, and life cycle involves self-fertilization, with the production of encysted eggs stored in the musculature of intermediate hosts.

3.4 Pathogenicity

Involves the disruption of skeletal musculature in respective intermediate and definitive hosts, with pronounced adverse in the latter.

3.5 Epidemiology and management

The infection is common in areas with high dependence on animal protein (beef, pork, fish), as well as intermittent contact with domestic and wild canids, especially in unhygienic conditions. Management involves good sanitary conditions of living and cooking, as well as minimal contacts with unhealthy canids.

4.0 Conclusion

They are efficiently adapted endoparasites, taking advantage of their well developed scolex (head) which is armed with suckers and hooks for

attachment. They also rely on their highly efficient integument armed with microvilli structures to absorb digested food within the gut.

5.0 Summary

- Cestodes are endoparasites of vertebrates
- Mode of transmission to human is through poorly cooked beef, pork or fish.
- In animals transmission is by faeco-oral means through consumption of cysts (eggs covered in protective casing).

6.0 Tutor-marked Assignment

Using annotated diagramme describe the life cycle of a named cestode.

7.0 References/Further reading

Yoloye, L.V. (1984) *Invertebrate Zoology*. Ilorin University Press, Ilorin. 318pp.

Hickman *et al.* (2008). *Integrated principles of Zoology*. 14th Edition, Mc-Graw Hill. 910pp.

Unit 2 Nematode

Content

- 1.0 Introduction
- 2.0. Objective
- 3.0. Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Life Cycle
 - 3.4 Pathogenicity
 - 3.5 Epidemiology and Management
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/Further reading

1.0 Introduction

Nematodes are round, cylindrical worms with pointed ends. they are ubiquitous in distribution, and are found parasitizing humans and wild life in different climes.

2.0. Objective

At the end of the class, student must be familiar with the different nematode classes, common nematode parasitic diseases, medical importance and chemotherapy.

3.0. Main content

3.1 Taxonomy:

There are basically two Classes namely Phasmidea and Aphasmeida, with the taxonomic character being the absence/presence of phasmids.

3.2 Biology (Characteristic Features)

Nematodes are round, elongate, cylindrical unsegmented worms. Integument is non-cellular, and musculature is of the longitudinal type. Nematodes are dioecious, with distinct sexually dimorphism. They are also characterized sensory papillae at both ends of the body (amphids). Examples include *Ascaris*, *Dracunculus*, *Loa*, *Onchocerca*, *Wuchereria*, *Brugia*, *Ancylostoma*, *Necator*, *Enterobius*, *trichinella*, and *Trichuris*.



A Soil transmitted nematode

3.3 Life Cycle

Sexes are separate, with copulation and fertilization occurring in the definitive hosts. Some have direct life cycle with re-infection enhanced by poor sanitary conditions, while others have indirect life cycle involving one or more intermediate hosts or vectors.

3.4 Pathogenicity

Involves impairment of tissues such as the skeletal musculature, subcutaneous tissues, GIT, respiratory system, as well as the eyes. Juvenile stages known as filarial worms of some members of this parasite groups are the cause of a number of disfiguring diseases which are prevalent in the tropics

These includes:

Onchocerciasis River blindness – cause by *Onchocerca volvulus* Guinea worm infection- caused by *Dracunculus medinensis* Filaris of all kinds

- a. Loasis (Calabar swelling)- caused by *Loa loa*
- b. Elephantiasis (bancroftian filariasis- caused by *Wuchereria bancrofti*

Hook worm- caused by *Necator americana* and *Ancylostoma deudenale*

In-Text Question

- What parasite causes river blindness?

Answer

- *Onchocerca volvulus*

3.5 Epidemiology and management

Involves vulnerable groups with high exposure to insanitary conditions (lack of pipe-borne water and decent toilet facilities). They are ubiquitous i.e widespread but very common in the tropics where temperature is optimal for their vectors. A good number of them are transmitted by arthropod vectors.

Management practices for nematode control includes; good sanitary

conditions, chemoprophylaxis and chemotherapy (Ivermectin, Mebendazole etc).

4.0 Conclusion

Nematodes constitute one of the largest collections of parasitic species infecting vertebrate hosts. They are often associated with poor sanitation which is prevalent in poor underdeveloped nations of the world.

5.0 Summary

Nematodes are called round worms because of their cylindrical appearance.

They have a global distribution and they comprise one of the largest parasite groups. They cause a number of debilitating and deforming diseases such as River blindness etc.

6.0 Tutor Marked Assignment (TMA)

- a. Make a list of different nematode parasites, their vectors and the disease they cause in humans.
- b. Draw the structure of the common round worm *Ascaris lumbricoides*

7.0 References/Further reading

- Hickman *et al.* (2008). Integrated principles of Zoology. 14th Edition, Mc-Graw Hill. 910pp
- Yoloye, L.V. (1984) *Invertebrate Zoology*. Ilorin University Press, Ilorin. 318pp.

Module 3


Unit 3 Acanthocephalans

- 1.0 Introduction
- 2.0. Objective
- 3.0. Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Life Cycle
 - 3.4 Pathogenicity
 - 3.5 Management
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/Further reading

Hickman *et al.* (2008). Integrated principles of Zoology. 14th Edition, Mc-Graw Hill.910pp.

1.0 Introduction

Acanthocephala (Greek ἄκανθος, *akanthos*, thorn + κεφαλή, *kephale*, head) is a phylum of parasitic worms known asacanthocephales, thorny-headed worms, or spiny-headed worms, characterized by the presence of an eversible proboscis, armed with spines, which it uses to pierce and hold the gut wall of its host. Acanthocephalans typically have complex life cycles, involving a number of hosts, including invertebrates, fishes, amphibians, birds, and mammals.



Structure of an Acanthocephalan

2.0. Objective

At the end of the class, student must be familiar with the parasitic acanthocephalans, their pathogenicity and epidemiology.

3.0. Main content

3.1 Taxonomy

About 1150 species have been described. The most common acanthocephalan includes *Macracanthorhynchus sp.*, the parasite of pigs.

3.2 Biology (Characteristic Features)


Acanthocephalans are highly adapted to a parasitic mode of life, and have lost many organs and structures through evolutionary processes. The size of the animals varies greatly, from forms a few millimetres in length to *Gigantorhynchus gigas*, which measures from 10 to 65 centimetres (3.9 to 26 in). The body surface of the acanthocephalan is peculiar. Externally, the skin has a thin cuticle covering the epidermis, which consists of a syncytium with no cell walls. The syncytium is traversed by a series of branching tubules containing fluid and is controlled by a few wandering, amoeboid nuclei. Inside the syncytium is an irregular layer of circular muscle fibres, and within this again some rather scattered longitudinal fibres; there is no endothelium. In their micro-structure the muscular fibres resemble those of nematodes.

Members are characterized by a cylindrical invaginated proboscis bearing rows of spines, hence the common name 'spiny-headed worms'. These spines are used for attachment to hosts. Members are endoparasites of vertebrates, especially fish, birds and mammals.

3.3 Life Cycle

Sexes are separate in acanthocephalans, with immature stages developing in invertebrates such as crustaceans and/or insects. Fully developed embryos are retained within the shell, egested with host faeces and remain unshelled until uptake by intermediate host. Acanthocephalans have complex life cycles, involving a number of hosts, for both developmental and resting stages. Complete life cycles have been worked out for only 25 species.

The lifecycle of *Polymorphus* spp., acanthocephalan parasites of birds.



In-Text Questions

- **Mention 4 possible hosts in the life cycle of Acanthocephalans**

Answer

- **Acanthocephalans typically have complex life cycles, involving a number of hosts, including invertebrates, fishes, amphibians, birds, and mammals.**

3.4 Pathogenicity

They are generally intestinal parasites, especially the duodenal arm of the small intestine. The nutritional requirement of these organisms, as well as release of metabolic wastes into host system severely impairs host, manifesting in clinical symptoms such as emaciation, anaemia and stunted growth and development.

3.5 Epidemiology and Management

They are cosmopolitan in distribution. Management procedure includes general hygienic condition, proper cooking of animal protein, chemotherapy.

4.0 Conclusion

Acanthocephalans are a bizarre and poorly studied parasitic group which are important causes of very disturbing gastric and intestinal conditions. Prevalence is common among poor and aqualor dwelling coastal communities.

6.0 Tutor Marked Assignment (TMA) 43

- a. Design a checklist of acanthocephalans of medical importance
- b. Describe a typical acanthocephalan life cycle.

7.0 **References/Further reading**

Hickman *et al.* (2008). Integrated principles of Zoology. 14th Edition, Mc-Graw Hill.910pp.

Module 3

Unit 4 Leeches

- 1.0 Introduction
- 2.0. Objective
- 3.0. Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Life Cycle
 - 3.4 Pathogenicity
- Epidemiology and Management
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor Marked Assignment (TMA)
- 7.0 References/Further reading

1.0 Introduction

Leeches, commonly called blood leeches are haematophagus, feeding on vertebrate host. They are aquatic and considered to be temporary ecto-parasites, separating from their hosts after blood meal. Leeches are segmented worms that belong to the phylum Annelida and comprise the subclass Hirudinea. Like other oligochaetes such as earthworms, leeches share a clitellum and are hermaphrodites. Leeches, such as the *Hirudo medicinalis*, have been historically used in medicine to remove blood from patients.

2.0. Objective

At the end of the class, student must be familiar with the *Hirudo species*, their habitat and mode of parasitism.

3.0 Main Content

3.1 Taxonomy

Leeches belong to the phylum Annelida and Class Hirudinea. A common example is *Hirudo sp*


In-Text Question

- **Mention one historical use of a named leech**
- **The *Hirudo medicinalis* have been historically used in medicine to remove blood from patients.** 45

3.2 Biology (Characteristic Features)

Leeches are dorso-ventrally flattened and segmented worms, with anterior and posterior suckers present for attachment to host and movement. Leeches are ectoparasitic, hence all organ systems are well developed. Their suckers are for attachment to their host during blood meal. Leeches are able to display a variety of behaviors that allow them to explore their environments and feed on their hosts. Exploratory behavior includes head movements and body waving

Most leeches do not feed on human blood, but instead prey on small invertebrates, which they eat whole. To feed on their hosts, leeches use their anterior suckers to connect to hosts for feeding, and also release an anesthetic to prevent the hosts from feeling them. Once attached, leeches use a combination of mucus and suction to stay attached and secrete an anticoagulant enzyme, hirudin, into the hosts' blood streams. Though certain species of leeches feed on blood, not all species can bite;



Dorsal and ventral view of the sucker of the leech, *H. medicinalis*

In-Text Question

- How do leeches feed on their hosts?

Answer

- To feed on their hosts, leeches use their anterior suckers to connect to hosts, and also release an anesthetic to prevent the hosts from feeling them. Once attached, they use a combination of mucus and suction to stay attached and secrete an anticoagulant enzyme, hirudin, into the hosts' blood streams

3.3 Life Cycle

Members are hermaphrodites, with exchange of gametes between two individuals, with eggs (in cocoons) buried in damp/moist or aquatic sites. Leeches reproduce by reciprocal fertilization, and sperm transfer occurs during copulation. Similar to the earthworms, leeches also use a clitellum to hold their eggs and secrete the cocoon.

3.4 Pathogenicity

Involves benign skin inflammation upon blood meal, as well enhance secondary infection via viral, bacterial and fungal agents.

3.5 Epidemiology and management

Leeches are found in fresh water bodies, particularly in slow moving streams, rivers, lakes and pond of the tropics, where residents have intermittent contact with water.

4.0 Conclusion

Leeches are a controversial parasitic group because most feed on small invertebrates by way of predation. Only a few feed on larger vertebrates by way of parasitism

5.0 Summary

- Leeches belong to the invertebrate class Annelida
- Like other oligochaetes such as earthworms, leeches share a clitellum and are hermaphrodites
- Only a few species live as ecto-parasites, the rest are predators.

6.0 Tutor Marked Assignment (TMA)

- a. Leeches are a controversial group of parasites. Discuss.

7.0 References/Further reading

Hickman *et al.* (2008). Integrated principles of Zoology. 14th Edition, Mc-Graw Hill. 910pp

Unit 5 Arthropods

Contents

- 1.0 Introduction
- 2.0 Objective
- 3.0 Main content
 - 3.1 Taxonomy
 - 3.2 Biology
 - 3.3 Life Cycle
 - 3.4 Pathogenicity
 - 3.5 Epidemiology and Management
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further reading

1.0 Introduction

Arthropods are winged invertebrates which are characterized by the possession of cuticular exo- skeleton. They vary widely in terms of their general body divisions, number of appendages, life cycle stages and habitat. They are the most successful animals on land and have a wide variety of niches, including those in which they act either as parasite vectors or parasites themselves (ecto-parasites).

2.0 Objective

At the end of the class, student must be the general arthropod taxonomy, the arthropod groups of parasitic importance (insect and acarina)

3.0 Main content

3.1 Taxonomy

Insects and Arachnids of parasitic importance. Quite a number of these organisms are of veterinary and medical importance, where they are considered vectors of diseases. Some of the important ones include ticks and mites, head and body lice, bedbugs, mosquitoes, tsetse flies, houseflies, deer flies, black flies, sand flies, biting midges.

3.2 Biology (Characteristic Features)

Members are characterized by the presence of jointed appendages, variable morphology, habitats and food types.

In-Text Question

- **Give examples of important arthropods**

Answer

- **Examples include ticks and mites, head and body lice, bedbugs, mosquitoes, tsetse flies, houseflies, deer flies, black flies, sand flies**

3.3 Life Cycle

This involves complete and/or incomplete metamorphosis. Arthropods with complete metamorphosis are usually more common as vectors of diseases, due to their varied diet and habitats, while those with incomplete metamorphosis are usually more common as parasites.

3.4 Pathogenicity

Many of these parasitic arthropods induce allergic reactions, dermatitis, some feed on subcutaneous tissues, as well as direct/indirect transmission of disease agents.

3.5 Epidemiology and Management

These arthropods are associated with age-groups in unhygienic environments.

Management: good sanitary conditions, chemotherapy

4.0 Conclusion

Arthropods are a unique group of animals because some groups are vectors (conveyors) of parasites, while others are themselves parasites (e.g. Ticks and Mites).

5.0 Summary

- Arthropods are the most successful animals on land.
- They have diverse forms of existence, one of which is by parasitism.
- Arthropod ecto-parasites are common in the Order Acarina
- Arthropods may serve as vectors either mechanically or biologically.

6.0 Tutor-Marked Assignment

a. Design a catalogue of arthropod species specimen of medical importance, stating their host and whether they are vectors or ecto-parasites.

7.0 References/Further reading

Hickman *et al.* (2008). *Integrated principles of Zoology*. 14th Edition, Mc-Graw Hill. 910pp

Yoloye, L.V. (1984) *Invertebrate Zoology*. Ilorin University Press, Ilorin. 318pp.